

Chlorphenamine Maleate Injection



Active Constituent

7mg/mL Chlorphenamine (equivalent to 10mg/mL Chlorphenamine Maleate)

Actions

Chlorpheniramine maleate is a first-generation antihistamine that antagonizes histamine H1 receptors. It is used for the symptomatic treatment of conditions associated with abnormal histamine activation, such as allergic reactions.

Indications

For the treatment of allergic reactions in horses, cattle, sheep, goats, pigs, dogs and cats.

Contraindications

Do not use in cases of known hypersensitivity to the product.

Precautions

The safety of the product has not been established during pregnancy and lactation.

Side Effects

May cause mild sedation.

Dosage & Administration

Use **subcutaneously** or **intramuscularly** at the following dose rates:

- Dogs, Cats, up to 10 kg bodyweight: 0.5 mL
- Dogs over 10 kg: 0.5 mL per 10 kg bodyweight
- Goats, Pigs, Sheep: 10-15 kg bodyweight: 1 - 2 mL.
15-25 kg bodyweight: 2 - 3 mL
- Cattle, Horses: 5 mL per 500 kg bodyweight

General Directions

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

Withholding Periods

WITHHOLDING PERIODS:

Meat (cattle, sheep, goats and pigs): Zero (0) Days

Milk (cattle, sheep and goats): Zero (0) Days

Any variation by the prescribing veterinarian to the approved use pattern may require an extended withholding period.

EXPORT SLAUGHTER INTERVAL (ESI): An ESI has not been established for this product. Note – observing the meat withholding period may not be sufficient to mitigate potential risks to export trade. Trade advice should be sought from Avet Health Ltd on 1300 28 38 28 before using this product.

First Aid

If poisoning occurs, contact a doctor or Poisons Information Centre on 13 1126 (Australia).

Safety Data Sheet

For Safety Data Sheet see www.avaxet.health

Presentation

50ml vial

Disposal

Dispose of used packaging by wrapping in paper and putting in garbage.

Storage

Store below 30 °C (room temperature). Following first broach, the vial must be stored at 2 °C to 8 °C (refrigerated). Do not freeze. Store upright. Protect from light.

Use the contents within 28 days of first broaching the vial

Poisons schedule

S4

Registration Numbers

APVMA Approval Number: 147379